

-
- 1 AKT phosphorylation associates with LOH of PTEN and leads to chemoresistance for gastric cancer.
Oki E., et al.
Int. J. Cancer

 - 2 Circulating cancer cells.
Oki E., et al.
Handbook of Immunohistochemistry and in situ hybridization Vol.4

 - 3 Genetic mutual relationship between PTEN and p53 in gastric cancer.
Oki E., et al.
Cancer Letters

 - 4 Impact of loss of heterozygosity of PTEN on the prognosis of gastric cancer.
Oki E., et al.
Gastroenterol. Hepatol.

 - 5 Overexpression of cyclooxygenase-2 and tumor angiogenesis in human gastric cancer.
Koga T., et al.
Hepato-Gastroenterol. 51(60):1626-30, 2004

 - 6 Analysis of gastric cancer with cDNA microarray.
Haraguchi N., et al.
Cancer Chemoth Pharm 54(S1):S21-S24, 2004

 - 7 Distribution of synchronous and metachronous multiple colorectal cancers.
Ikeda Y., et al.
Hepato-Gastroenterol. 51(56):443-446, 2004

 - 8 Effective follow-up for recurrence or a second primary cancer in patients with early gastric cancer.
Ikeda Y., et al.
Br. J. Surg.

 - 9 Prophylactic lymph node dissection for early gastric cancer invading submucosa.
Ikeda Y., et al.
Hepato-Gastroenterol. 51(57):887-890, 2004
-

-
- 10 *p53* mutation profiling of multiple esophageal carcinoma using laser capture microdissection to demonstrate field carcinogenesis.
-
- Ito S., et al.
-
- Int. J. Cancer 113(1):22-28, 2005
-
- 11 A variant form of *hMTH1*, a human homologue of *E. coli mutT* gene, correlates with somatic mutation in *p53* tumour suppressor gene in gastric cancer patients.
-
- Kimura Y., Oda S., Egashira A., Kakeji Y., Baba H., Nakabeppu Y., Maehara Y.
-
- J Med Genet 41:e57, 2004
-
- 12 Detection of occult cancer cells in peripheral blood and bone marrow by quantitative RT-PCR assay for cytokeratin-7 in breast cancer patients.
-
- Masuda T., et al.
-
- Int. J. Oncol.
-
- 13 Common fragile genes.
-
- Matsuyama A., et al.
-
- Eur. J. Histochem. 48(1):24-30, 2004
-
- 14 Clinical significance of enhancer of zeste homolog 2 expression in colorectal cancer cases.
-
- Mimori K., et al.
-
- Eur. J. Surg. Oncol.
-
- 15 Clinical significance of the overexpression of the candidate oncogene *CYP24* in esophageal cancer.
-
- Mimori K., et al.
-
- Ann. Oncol. 15:236-241, 2004
-
- 16 Coexpression of matrix metalloproteinase-7 (MMP-7) and epidermal growth factor (EGF) receptor in colorectal cancer: An EGF receptor tyrosine kinase inhibitor is effective against MMP-7-expressing cancer cells.
-
- Mimori K., et al.
-
- Clin. Cancer Res. 10:8243-8249, 2004
-
- 17 Identification of molecular markers for metastasis-related genes in primary breast cancer cells.
-
- Mimori K., et al.
-
- Clin. Exp. Metastas.
-

-
- 18 Prediction of 5'-deoxy-5-fluorouridine sensitivity in colorectal cancer cases by thymidine phosphorylase activity and preliminary identification of susceptibility related genes.

Mimori K., et al.

Oncol. Rep. 12:19-23, 2004

-
- 19 Loss of phenotypic expression is related to tumor progression in early gastric differentiated adenocarcinoma.

Nakamura T., et al.

Histopathology

-
- 20 Clinical significance in the preoperative measurement of serum value of lymphocyte subpopulation in patients with squamous cell carcinoma of the esophagus.

Nozoe T., et al.

World J. Gastroenterol.

-
- 21 Clinicopathologic significance of cyclin A expression in colorectal carcinoma.

Nozoe T., et al.

J. Exp. Clin. Cancer Res. 23:127-133, 2004

-
- 22 Correlation of ABO blood group with clinicopathologic characteristics of patients with esophageal squamous cell carcinoma.

Nozoe T., et al.

Dis Esophagus 17:146-149, 2004

-
- 23 Correlation of p53 over-expression with gender of patients with squamous cell carcinoma of the esophagus.

Nozoe T., et al.

Hepato-Gastroenterol.

-
- 24 Correlation of pre-operative coexistence of organ dysfunction with overall survival of patients with squamous cell carcinoma of the thoracic esophagus.

Nozoe T., et al.

Oncol. Rep. 12:655-658, 2004

-
- 25 Relationship between intracellular localization of p34cdc2 protein and differentiation of esophageal squamous cell carcinoma.
-
- Nozoe T., et al.
-
- J. Cancer Res Clin Oncol
-
- 26 Significance of immunohistochemical over-expression of CD44v6 as an indicator of malignant potential in esophageal squamous cell carcinoma.
-
- Nozoe T., et al.
-
- J. Cancer Res Clin Oncol 130:334-338, 2004
-
- 27 Significance of immunohistochemical over-expression of cyclooxygenase-2 in squamous cell carcinoma of the esophagus.
-
- Nozoe T., et al.
-
- Am. J. Surg.
-
- 28 Microsatellite Instability in Gastrointestinal Tract Cancers: A Brief Update.
-
- Oda S., et al.
-
- Surg. Today
-
- 29 Microsatellite instability in human cancers: Its virtual and real images.
-
- Oda S., et al.
-
- Dynamical Genetics 167-184, 2004
-
- 30 The high expression of Fractalkine results in a better prognosis for colorectal cancer patients.
-
- Ohta M., et al.
-
- Int. J. Oncol. 26:41-47, 2005
-
- 31 A case of gastric hyperplastic polyp with malignant transformation.
-
- Shibahara K., et al.
-
- Hepato-Gastroenterol.
-
- 32 Targeted disruption of one allele of the *Y-box binding protein-1(YB-1)* gene in mouse embryonic stem cells and increased sensitivity to cisplatin and mitomycin C.
-
- Shibahara K., et al.
-
- Cancer Sci. 95(4):348-353, 2004
-

-
- 33 Increased antitumor activity in combined treatment TS-1 and docetaxel ~ a preclinical study using gastric cancer xenografts.
-
- Takahashi I., et al.
-
- Oncology
-
- 34 End- to-end anastomosis using a surgical instrument after a segmental resection of the distal duodenum.
-
- Toh Y., Yamamoto M., Endo K., Ikeda Y., Baba H., Kohnoe S., Okamura T.
-
- J Surg Oncol 85:88-90, 2004
-
- 35 Expression of the metastasis-associated MTA1 protein and its relationship to deacetylation of the histone H4 in esophageal squamous cell carcinomas.
-
- Toh Y., et al.
-
- Int. J. Cancer 110(3):362-367, 2004
-
- 36 Neoangiogenesis in patients with gastric carcinoma in relation to the expression of vascular endothelial growth factor and thymidine phosphorylase.
-
- Tsujitani S., et al.
-
- Anticancer Res. 24(3b):1853-1859, 2004
-
- 37 Identification of HLA-A24-Restricted CTL epitope from cancer-testis antigen, NY-ESO-1, and induction of a specific antitumor immune response.
-
- Yamaguchi H., et al.
-
- Clin. Cancer Res. 10:890-896, 2004
-
- 38 Distributions in CEA doubling time differ in patients with recurrent colorectal carcinomas.
-
- Yamamoto M., Maehara Y., Sakaguchi Y., Mine H., Yamanaka T., Korenaga D., Okamura T.
-
- Hepato-Gastroenterology 51(55):147-151, 2004
-
- 39 Polyps with different grades of dysplasia and their distribution in the colorectum.
-
- Yamamoto M., Mine H., Kusumoto H., Maehara Y., Sugimachi K.
-
- Hepato-Gastroenterology 51(55):121-123, 2004
-

-
- 40 Postoperative morbidity / mortality and survival rates after total gastrectomy, with splenectomy/pancreaticosplenectomy for patients with advanced gastric cancer.

Yamamoto M., Baba H., Kakeji Y., Endo K., Ikeda Y., Toh Y., Kohnoe S., Okamura T., Maehara Y.

Hepato-Gastroenterology 51(55):298-302, 2004

-
- 41 Prognostic significance of tumor markers in peritoneal lavage in advanced gastric cancer.

Yamamoto M., Baba H., Kakeji Y., Endo K., Ikeda Y., Toh Y., Kohnoe S., Okamura T., Maehara Y.

Oncology 67(1):19-26, 2004

-
- 42 N-cadherin is regulated by activin A and associated with tumor aggressiveness in esophageal carcinoma.

Yoshinaga K., Inoue H., et al.

Clin. Cancer Res. 10(17):5702-07, 2004

1 Granulomatous Lobular Mastitis Misdiagnosed as Breast Carcinoma

Tokunaga E.,Kimura Y.,Kitamura K.,Maehara Y.

Breast J 10(3):261-262, 2004

2 No hypersensitive estrogen receptor- mutation (K303R) in Japanese breast carcinomas

Tokunaga E.,Kimura Y.,Maehara Y.

Breast Cancer Research and Treatment 84:289-292, 2004

3 Sensitization of chemotherapy by anti-HER.

Kataoka A., et al.

Breast Cancer 11(2):105-115, 2004

4 Team approach to providing the multidisciplinary medical treatment derived by the patients and their family.

Kataoka A., et al.

Breast Cancer

5 Brain metastasis of breast cancer.

Ohno S., et al.

Breast Cancer 11(1):27-29, 2004

6 Clinical significance of Caveolin-1, Caveolin-2 and HER2/neu mRNA expression in human breast cancer.

Sagara Y., et al.

Br. J. Cancer 91:959-965, 2004

7 Clinicopathological feature and long-term prognosis of apocrine carcinoma of the breast in Japanese women.

Takeuchi H., et al.

Breast Cancer Res. Treat. 88(1):49-54, 2004

8 Surgical results of total pelvic exenteration for locally advanced colorectal adenocarcinoma.

Takeuchi H., et al.

Hepato-Gastroenterol.

-
- 9 Prognostic implication of p53 protein expression in relation to nuclear pleomorphism and the MIB-1 counts in breast cancer.

Tsutsui S., et al.

Breast Cancer 11(2):160-168, 2004

-
- 10 The reduced expression of PTEN protein and its prognostic implications in invasive ductal carcinoma of the breast.

Tsutsui S., et al.

Oncology

-
- 1 Long-term function of cryopreserved rat hepatocytes in a coculture system.

Sugimachi K., et al.

Cell Transplant. 13(2):187-195, 2004

-
- 2 cDNA microarray analysis in hepatocyte differentiation in Huh 7 cells.

Yamashita Y., et al.

Cell Transplant. 13:793-799, 2004

-
- 3 Muscle-targeted interleukin-12 gene therapy of orthotopic hepatocellular carcinoma in mice using in vivo electrosonoporation.

Yamashita Y., Shimada M., Minagawa R., Tsujita E., Harimoto N., Tanaka S., Shirabe K., Miyazaki J., Maehara Y.

Mol Cancer Ther. 3(9):1177-82, 2004

-
- 4 Combined treatment of histone deacetylase inhibitor with PPAR-gamma ligand enhances the liver specific function of HepG2.

Tsujita E., et al.

Cell Transplant.

-
- 5 Hepatobiliary and pancreatic: peritoneal dissemination mimicking liver metastases.

Harimoto N., et al.

J. Gastroenterol. Hepatol. 19:938, 2004

-
- 6 Images of interest. Hepatobiliary and pancreatic: peritoneal dissemination mimicking liver metastases.

Harimoto N., Shimada M., Shirabe K., Tanaka S., Taketomi A., Tsujita E., Maehara S., Itoh S., Kitagawa D., Honda H., Maehara Y.

J Gastroenterol Hepatol. 19(8):938, 2004

-
- 7 The newly established human hepatocyte cell line: application for the bioartificial liver.

Harimoto N., et al.

J. Hepatol.

-
- 8 The role of heat shock protein 27 expression in hepatocellular carcinoma in Japan: special reference to the difference between hepatitis B and C.

Harimoto N., Shimada M., Aishima S., Kitagawa D., Itoh S., Tsujita E., Maehara S., Taketomi A., Tanaka S., Shirabe K., Maehara Y.

Liver Int. 24(4):316-321, 2004

-
- 9 Aberrant Expression of Laminin Gamma 2 Chain and Its Prognostic Significance in Intrahepatic Cholangiocarcinoma According to Growth Morphology.

Aishima S., Matsuura S., Terashi T., Taguchi K., Shimada M., Maehara Y., Tsuneyoshi M.

Modern Pathology 17:938-945, 2004

-
- 10 Role of Expression of Focal Adhesion Kinase on Progression in Hepatocellular Carcinoma.

Itoh S., Maeda T., Shimada M., Aishima S., Shirabe K., Tanaka S., Maehara Y.

Clinical Cancer Research 10:2812-2817, 2004

-
- 11 Clinicopathological correlates of aspartyl (asparaginy) -hydroxylase over-expression in cholangiocarcinoma

Maeda T., Taguchi K., Aishima S., Shimada M., Hintz D., LaRusso N., Gores G., Tsuneyoshi M., Sugimachi K., Wands JR., Monte S.

Cancer Detection and Prevention 28:313-318, 2004

-
- 12 Role of tissue trace elements in liver cancers and non-cancerous liver parenchyma.

Maeda T., et al.

Hepato-Gastroenterol.

-
- 13 Selenoprotein P, as a predictor for evaluating gemcitabine resistance in human pancreatic cancer cells.

Maehara S., Tanaka S., Shimada M., Shirabe K., Saito Y., Takahashi K., Maehara Y.

International Journal of Cancer 112:184-189, 2004

-
- 14 Prognostic factors in node-negative intrahepatic cholangiocarcinoma with special reference to angiogenesis.

Shirabe K., Shimada M., Tsujita E., Aishima S., Maehara S., Tanaka S., Takenaka K., Maehara Y.

Am J Surg. 187:538-542, 2004

-
- 15 A novel intrahepatic arterial chemotherapy after radical resection for advanced hepatocellular carcinoma.

Tanaka S., et al.

Hepato-Gastroenterol.

-
- 16 CCN4 and CCN6 variants in Wnt-inducible signaling pathway. in The CCN proteins: A new family of cell growth and differentiation regulators.

Tanaka S., et al.

Imperial College Press, London

-
- 17 Surgical outcome of patients with hepatocellular carcinoma originating in the caudate lobe.

Tanaka S., et al.

Am. J. Surg.

-
- 18 Surgical radiofrequency ablation for treatment of hepatocellular carcinoma: an endoscopic or open approach.

Tanaka S., et al.

Hepato-Gastroenterol.

-
- 19 A gene-expression signature can quantify the degree of hepatic fibrosis in the rat.

Utsunomiya T., et al.

J. Hepatol. 41:399-406, 2004

-
- 20 Clinical significance of elongation factor-1 delta mRNA expression in oesophageal carcinoma.

Utsunomiya T., et al.

Br. J. Cancer 91:282-286, 2004

-
- 21 Clinical significance of human kallikrein gene 6 (*KLK6*) mRNA expression in colorectal cancer.

Utsunomiya T., et al.

Clin. Cancer Res.

-
- 22 Clinicopathologic and prognostic values of apolipoprotein D alterations in hepatocellular carcinoma.

Utsunomiya T., et al.

Int. J. Cancer

23 Expression of cancer-testis antigen (CTA) genes in intrahepatic cholangiocarcinoma.

Utsunomiya T., et al.

Ann. Surg. Oncol. 11(10):934-940, 2004

-
- 1 Preoperatively undiagnosable intrahepatic bile duct dilatation contains early-stage intrahepatic cholangiocarcinoma.

Kinjo N., et al.

Hepato-Gastroenterol.

-
- 2 Propranolol alone may not be acceptable to prevent first esophageal variceal bleeding in Japanese cirrhotic patients: randomized controlled trial.

Tomikawa M., et al.

J. Gastroenterol. Hepatol. 19(5):576-581, 2004

-
- 3 Essential role of PDGFR α -p70S6K signaling in mesenchymal cells during therapeutic and tumor angiogenesis in vivo.

Tsutsumi N., et al.

Circulation Res. 94(9):1186-94, 2004

-
- 1 Portal vein thrombosis complicated with disseminated intravascular coagulation due to acute cholecystitis.

Inoguchi H., et al.

Hepato-Gastroenterol. 51(57):661-663, 2004

-
- 2 Hypereosinophilic syndrome accompanying gangrene of the toes with peripheral arterial occlusion.

Funahashi S., et al.

Angiology

-
- 3 Pedal artery bypass for diabetic arteriosclerosis in the bilateral lower limbs: report of a case.

Yamaoka T., et al.

Breakthrough in the treatment for critical limb ischemia in Japan 97-100, 2004

-
- 1 Loss of heterozygosity (LOH) in non-small cell lungcancer: difference between adenocarcinoma and squamous cell carcinoma.

Yoshino I., Osoegawa A., Yohena T., Kameyama T., Oki E., Oda S., Maehara Y.

Respiratory Medicine

-
- 2 Multimodal treatment for resectable epithelial type malignant pleural mesothelioma.

Yoshino I., Yamaguchi M., Okamoto T., Ushijima C., Fukuyama Y., Ichinose Y., Maehara Y.

World Journal of Surgical Oncology May 52(1):11, 2004

-
- 3 Sleeve tracheal resection through an anterior approach.

Yoshino I., et al.

Asian Thoracic Cardiovasc Ann.

-
- 4 Preoperative mechanical bowel preparation unnecessary in patients undergoing thoracic surgery.

Yamazaki K., et al.

Jpn. J. Thorac. Cardiovasc. Surg. 52(9):407-410, 2004

-
- 5 Identification of MGB1 as a marker in the differential diagnosis of lung tumors in patients with a history of breast cancer by analysis of publicly available SAGE data.

Koga T., et al.

J. Mol. Diagn. 6(2):90-95, 2004

-
- 6 Hepatocellular carcinoma in a case of Dubin-Johnson syndrome treated successfully with a central bilateral segmentectomy.

Shikada Y., Matsumata T., Suehiro T., Sugimachi K.

Hepatogastroenterology 51:833-836, 2004

-
- 7 V 1+ T cells producing CC chemokines may bridge a gap between neutrophils and macrophages in innate immunity during *Escherichia coli* infection in mice

Tagawa T., Nishimura H., Yajima T., Hara H., Kishihara K., Matsuzaki G., Yoshino I., Maehara Y., Yoshikai Y.

Journal of Immunology 173:5156-5164, 2004

-
- 8 Hypermethylation of FHIT as a prognostic marker in nonsmall cell lung carcinoma.

Maruyama R., et al.

Cancer 100(7):1472-77, 2004

-
- 9 Prognostic value of visceral pleural invasion in resected non-small cell lung cancer diagnosed by using a jet stream of saline solution.
-
- Maruyama R., et al.
-
- J. Thorac. Cardiovasc. Surg. 127(6):1587-92
-
- 10 Long-term survival after surgical resection of liver metastasis from lung cancer.
-
- Nagashima A., et al.
-
- Jpn. J. Thorac. Cardiovasc. Surg. 52(6):311-313, 2004
-
- 11 The significance of serum active matrix metalloproteinase 9 in patients with non-small cell lung cancer
-
- Ondo K., Sugio K., Yamazaki K., Yamaguchi M., Yano T., Yoshino I., Maehara Y.
-
- Lung Cancer 46:205-213, 2004
-
- 12 Regulation of p27 by Skp2 Is Associated With Aggressiveness In Non-Small-Cell Lung Cancer.
-
- Osoegawa A., Yoshino I., Tanaka S., Sugio K., Kameyama T., Yamaguchi M., Maehara Y.
-
- J Clin Oncol 22:4165-4173, 2004
-
- 13 Clinical significance of serum TIMP-1 and TIMP-2 in patients with non-small cell lung cancer.
-
- Suemitsu R., Yoshino I., Tomiyasu M., Fukuyama S., Okamoto T., Maehara Y.
-
- Surg Today 34:896-901, 2004
-
- 14 Serum tissue inhibitors of metalloproteinase-1 and -2 in patients with non-small cell lung cancer.
-
- Suemitsu R., et al.
-
- Surg. Today 34(11):896-901, 2004
-
- 15 The effects of pretreatment with donor antigen and immunosuppressive agents on fully allogenic tracheal graft.
-
- Suemitsu R., Yoshino I., Shoji F., Yamaguchi M., Tomita Y., Maehara Y.
-
- J Surg Res 122:8-13, 2004
-
- 16 Video-assisted extended thymectomy in patients with thymoma by lifting the sternum: Is it safe?
-
- Takeo S., et al.
-
- Ann. Thorac. Surg. 77(5):1878, 2004
-

-
- 17 Video-assisted thoracoscopic resection of a giant anterior mediastinal tumor (thymolipoma) using an original sternum-lifting technique.

Takeo S., et al.

Jpn. J. Thorac. Cardiovasc. Surg.

-
- 18 Elective video-assisted thoracoscopic lung biopsy for interstitial lung disease.

Yamaguchi M., Yoshino I., Suemitsu R., Osoegawa A., Kameyama T., Tagawa T., Fukuyama S., Maehara Y.

Asian Cardiovasc Thorac Ann 12:65-68, 2004

-
- 19 Surgical treatment of neurogenic tumors of the chest.

Yamaguchi M., Yoshino I., Fukuyama S., Osoegawa A., Kameyama T., Tagawa T., Maehara Y.

Ann Thorac Cardiovasc Surg 10:148-151, 2004

-
- 1 Involvement of Rho-kinase in cold ischemia-reperfusion injury after liver transplantation in rats.

Shiotani S., Shimada M., Suehiro T., Soejima Y., Yosizumi T., Shimokawa H., Maehara Y.

Transplantation 78(3):375-82, 2004

-
- 2 A simple and accurate formula to estimate left hepatic graft volume in living donor adult liver transplantation.

Harada N., Shimada S., Yoshizumi T., Suehiro T., Soejima Y., Maehara Y.

Transplantation 77(10):1571-5, 2004

-
- 3 IL-12 gene therapy is an effective therapeutic strategy for hepatocellular carcinoma in immunosuppressed mice.

Harada N., Shimada S., Okano S., Suehiro T., Soejima Y., Tomita Y., Maehara Y.

The journal of immunology 173(11):6635-44, 2004

-
- 4 Impact of preoperative serum C-reactive protein on the prognosis in patients with hepatocellular carcinoma.

Hashimoto K., et al.

Cancer

-
- 5 Granulocyte-colony stimulating factor producing gallbladder cancer without recurrence more than 2 year after resection.

Ikeda T., et al.

Surg. Today

-
- 6 Intestinal obstruction because of a migrated metallic biliary stent.

Ikeda T., et al.

Gastrointest. Endosc. 60(6):988-989, 2004

-
- 7 Limited hepatic resection for hepatocellular carcinoma in the caudate lobe.

Ikegami T., et al.

World J. Surg. 28:697-701, 2004

-
- 8 Neurilemmoma of the diaphragm mimicking a liver tumor: case report.

Ikegami T., et al.

Abdom. Imaging. 29:85-86, 2004

-
- 9 Primary gas gangrene of the pancreas: Report of a case.
-
- Ikegami T., et al.
-
- Surg. Today 34:80-81, 2004
-
- 10 Retractable mesenteritis: Report of a case.
-
- Ikegami T., et al.
-
- Surg. Today 34:547-549, 2004
-
- 11 Personal experience of laparoscopic cholecystectomy.
-
- Kakizoe S., et al.
-
- Hepato-Gastroenterol. 51:934-936, 2004
-
- 12 The critical role of fas-fas ligand interaction in donor-specific transfusion-induced tolerance to H-Y antigen.
-
- Minagawa R., et al.
-
- Transplantation 78(6):799-806, 2004
-
- 13 Characteristic gene expression induced by polyurethane foam/spheroid culture of hepatoma cell line, hep G2 as a promising cell source for bioartificial liver.
-
- Shimada M., et al.
-
- Hepato-Gastroenterol.
-
- 14 Des-gamma-carboxy prothrombin is a useful predictor of early recurrence of hepatocellular carcinoma after living donor adult liver transplantation.
-
- Shimada M., et al.
-
- Transplant P.
-
- 15 Is graft size a major risk factor in living-donor adult liver transplantation?
-
- Shimada M., Ijichi H., Yonemura Y., Harada N., Shiotani S., Ninomiya M., Yoshizumi T., Soejima Y., Suehiro T., Maehara Y.
-
- Transpl Int. 17:310-6, 2004
-

-
- 16 The impact of splenectomy or splenic artery ligation on the outcome of a living donor adult liver transplantation using a left lobe graft.

Shimada M., Ijichi H., Yonemura Y., Harada N., Shiotani S., Ninomiya M., Terashi T., Yoshizumi T., Soejima Y., Suehiro T., Maehara Y.

Hepatogastroenterology 51:625-9, 2004

-
- 17 Graft-versus-host disease following living donor liver transplantation.

Soejima Y., Shimada M., Suehiro T., Hiroshige S., Gondo H., Takami A., Yasue S., Maehara Y.

Liver Transplantation 3:460-464, 2004

-
- 18 Accelerated rehabilitation with early postoperative oral feeding following gastrectomy.

Suehiro T., et al.

Hepato-Gastroenterol. 51:1852-1855, 2004

-
- 19 Change in alpha glutathione s-transferase levels during liver resection.

Suehiro T., et al.

Hepato-Gastroenterol. 51:1747-1750, 2004

-
- 20 Preoperative hyaluronic acid measurement to assess the hepatic functional reserve.

Suehiro T., et al.

Hepato-Gastroenterol. 51:1776-1779, 2004

-
- 21 Decreased expression of osteopontin is related to tumor aggressiveness and clinical outcome of intrahepatic cholangiocarcinoma.

Terashi T., Aishima S., Taguchi K., Asayama Y., Sugimachi K., Matsuura S., Shimada M., Maehara S., Maehara Y., Tsuneyoshi M.

Liver Int. 24:38-45, 2004

-
- 1 Global analysis of altered gene expressions during the process of esophageal squamous cell carcinogenesis in the rat: a study combined with a laser microdissection and a cDNA microarray.

Nishida K., et al.

Cancer Res.

-
- 2 Liver metastasis from thyroid carcinoma 32 years after resection of the primary tumor: report of a case.

Kouso H., et al.

Surg. Today
